arXiv:1609.00839v1 [math.GM] 3 Sep 2016

How we pass from semigroups to hypersemigroups

Niovi Kehayopulu

Abstract. The aim of writing this paper is given in the title. The results on semigroups can be easily transferred to hypersemigroups in the way indicated in the present paper.

1 Introduction and prerequisites

An *hypergroupoid* is a nonempty set H with an hyperoperation

$$\circ : H \times H \to \mathcal{P}^*(H) \mid (a, b) \to a \circ b$$

on H and an operation

$$* : \mathcal{P}^*(H) \times \mathcal{P}^*(H) \to \mathcal{P}^*(H) \mid (A, B) \to A * B$$

on $\mathcal{P}^*(H)$ (induced by the operation of H) such that

$$A * B = \bigcup_{(a,b) \in A \times B} (a \circ b)$$

for every $A, B \in \mathcal{P}^*(H)$ ($\mathcal{P}^*(H)$ denotes the set of nonempty subsets of H). As the operation "$*$" depends on the hyperoperation "$\circ$", an hypergroupoid can be also denoted by $(H, \circ)$ (instead of $(H, \circ, *)$). If H is an hypergroupoid then, for every $x, y \in H$, we have $\{x\} * \{y\} = x \circ y$. A nonempty subset A of an hypergroupoid $(H, \circ)$ is called a *left* (resp. *right*) *ideal* of H if $H * A \subseteq A$ (resp. $A * H \subseteq A$). A subset of H which is both a left ideal and a right ideal of H is called an *ideal* of H. A nonempty subset A of H is called a *subsemigroup* of H if $A * A \subseteq A$. Clearly, every left (resp. right) ideal of H is a subsemigroup of H. An hypergroupoid H is called *hypersemigroup* if $\{x\} * (y \circ z) = (x \circ y) * \{z\}$ for every $x, y, z \in H$. This is equivalent to saying that $\{x\} * \Big(\{y\} * \{z\}\Big) = \Big(\{x\} * \{y\}\Big) * \{z\}$ for every $x, y, z \in H$. Identifying the $\{x\}$ by x and the $\{z\}$ by z we could write, for short, $x * (y \circ z) = (x \circ y) * z$ for all $x, y, z \in H$. If H is an hypersemigroup

and $A, B, C \in \mathcal{P}^*(H)$, then we have

$$\begin{aligned}(A * B) * C &= \bigcup_{(a,b,c)\in A\times B\times C} \Big((a \circ b) * \{c\}\Big) \\ &= \bigcup_{(a,b,c)\in A\times B\times C} \Big(\{a\} * (b \circ c)\Big) = A * (B * C) \\ &= \bigcup_{(a,b,c)\in A\times B\times C} \Big(\{a\} * \{b\} * \{c\}\Big).\end{aligned}$$

Thus we can write $(A * B) * C = A * (B * C) = A * B * C$.
From this, using induction, follows that for any finite family $A_1, A_2, ..., A_n$ of elements of $\mathcal{P}^*(H)$, we have

$$A_1 * A_2 * ..... * A_n = \bigcup_{(a_1,a_2...a_n)\in A_1\times A_2\times ...\times A_n} \Big(\{a_1\} * \{a_2\}*, ... * \{a_n\}\Big).$$

The following proposition, though clear, plays an essential role in the theory of hypergroupoids.

Proposition 1.1. *Let $(H, \circ)$ be an hypergroupoid, $x \in H$ and $A, B \in \mathcal{P}^*(H)$. Then we have the following:*

(1) $x \in A * B \iff x \in a \circ b$ *for some* $a \in A$, $b \in B$.

(2) *If* $a \in A$ *and* $b \in B$, *then* $a \circ b \subseteq A * B$.

Following Zadeh, any mapping $f : H \to [0, 1]$ of an hypergroupoid H into the closed interval $[0, 1]$ of real numbers is called a *fuzzy subset* of H (or a *fuzzy set* in H) and f_A (: the characteristic function of A) is the mapping

$$f_A : H \to [0, 1] \mid x \to f_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A. \end{cases}$$

The concepts of fuzzy left and fuzzy right ideals of semigroups can be naturally transferred in case of an hypergroupoid as follows: Let H be an hypergroupoid. A fuzzy subset f of H is called a *fuzzy left ideal* of H if $f(x \circ y) \geq f(y)$ for all $x, y \in H$, in the sense that $x, y \in H$ and $u \in x \circ y$ implies $f(u) \geq f(y)$. A fuzzy subset f of H is called a *fuzzy right ideal* of H if $f(x \circ y) \geq f(x)$ for all $x, y \in H$, meaning that if $x, y \in H$ and $u \in x \circ y$, then $f(u) \geq f(x)$. A fuzzy subset f of H is called a *fuzzy ideal* of H it is both a fuzzy left and a fuzzy right ideal of H. As one can easily see, a fuzzy subset f of H is a fuzzy ideal

of H if and only $f(x \circ y) \geq \max\{f(x), f(y)\}$ for all $x, y \in H$, in the sense that $x, y \in H$ and $u \in x \circ y$ implies $f(u) \geq \max\{f(x), f(y)\}$.

Fuzzy interior ideals of semigroups (without order) and fuzzy simple semigroups (without order) have been first considered by Kuroki in [2]. If S is a regular or an intra-regular ordered semigroup, then the ideals and the interior ideals of S coincide. In particular, if S is a *poe*-semigroup (i.e. an ordered semigroup having a greatest element with respect to the order), then the ideal elements and the interior ideal elements of S coincide as well. In an attempt to show the similarity between the theory of ordered semigroups and the theory of fuzzy ordered semigroups, we have proved in [1] that in regular and in intra-regular ordered semigroups the fuzzy ideals and the fuzzy interior ideals coincide. We also proved that a set A is an interior ideal of an ordered semigroup S if and only its characteristic function f_A is a fuzzy interior ideal of S. We finally proved that an ordered semigroup is simple if and only if it is fuzzy simple, equivalently, if every fuzzy interior ideal of S is a constant function. The present paper is based on the papers [1] and [2] in the References, and its aim is to show the way we pass from semigroups to hypersemigroups. Analogous results hold for ordered hypersemigroups as well. Results on semigroups can be translated into the language of the hypersemigroups in the same way. Although we cannot say "all the results", up today, I have never seen a result that cannot be translated. And this is why, when we work on an hypersemigroup, we should indicate inside the paper the 1–3 papers on semigroups on which our paper is based. Instead, the corresponding papers on semigroups are sometimes (not always) cited in the References among many other papers on hypersemigroups (or even on semigroups) not related with the investigation. Concerning the present paper, as one can see, only the papers on semigroups cited in the References are needed in the investigation and nothing else. We use the terms left (right) ideal, bi-ideal, quasi-ideal instead of left (right) hyperideal, bi-hyperideal, quasi-hyperideal and so on, and this is because in this structure there are no two kind of left ideals, for example, to distinguish them as left ideal and left hyperideal. The left ideal in this structure is that one which corresponds to the left ideal of semigroups. We will give further interesting information about the hypersemigroups (without order) in a forthcoming paper.

2 Main results

Definition 2.1. Let H be an hypersemigroup. A nonempty subset A of H is called an *interior ideal* of H if

$$H * A * H \subseteq A.$$

By a *subidempotent interior ideal* of H we mean an interior ideal of H which is at the same time a subsemigroup of H.

Definition 2.2. Let H be an hypergroupoid. A nonempty subset A of H is called a *fuzzy interior ideal* of H if

$$f\Big((x \circ a) * \{y\}\Big) \geq f(a) \text{ for every } x, a, y \in H.$$

For an hypersemigroup, we clearly have

$$(x \circ a) * \{y\} = \{x\} * (a \circ y) = \{x\} * \{a\} * \{x\}.$$

Proposition 2.3. *Let H be an hypersemigroup. If A is an interior ideal of H, then the characteristic function f_A is a fuzzy interior ideal of H. "Conversely", if A is a nonempty subset of H such that f_A is a fuzzy interior ideal of H, then A is an interior ideal of H.*

Proof. $\Longrightarrow$. Let $x, a, y \in H$. Then $f_A\Big((x \circ a) * \{y\}\Big) \geq f_A(a)$. In fact: Let $u \in (x \circ a) * \{y\}$. If $a \in A$, then $f_A(a) = 1$. Since A is an interior ideal of H, we have $H * A * H \subseteq A$. So we have $u \in \{x\} * \{a\} * \{y\} \subseteq H * A * H \subseteq A$. Then $u \in A$, and $f_A(u) = 1$. Thus we get $f_A(u) \geq f_A(a)$. Let now $a \notin A$. Then $f_A(a) = 0$. Since f_A is a fuzzy subset of H and $u \in H$, we have $f_A(u) \geq 0$. Thus we have $f_A(u) \geq f_A(a)$.
$\Longleftarrow$. Let A be a nonempty subset of H and f_A a fuzzy interior ideal of H. Then $H * A * H \subseteq A$. Indeed: Let $u \in H * A * H$. Then $u \in (x \circ a) * \{y\}$ for some $x, y \in H$, $a \in A$. Since f_A a fuzzy interior ideal of H, we have $f_A(u) \geq f_A(a) = 1$. Since f_A is a fuzzy subset of H and $u \in H$, we have $f_A(u) \leq 1$. So we have $f_A(u) = 1$, and $u \in A$. □

Proposition 2.4. *Let H be an hypersemigroup. If f is a fuzzy ideal of H, then f is a fuzzy interior ideal of H.*

Proof. Let $x, a, y \in H$. Then $f\Big((x \circ a) * \{y\}\Big) \geq f(a)$. In fact: Let $u \in (x \circ a) * \{y\}$. By Proposition 1.1, there exists $v \in x \circ a$ such that $u \in v \circ y$. Since f is a fuzzy left ideal of H, we have $f(x \circ a) \geq f(a)$ and, since $v \in x \circ a$, we get $f(v) \geq f(a)$. Since f is a fuzzy right ideal of H, we have $f(v \circ y) \geq f(v)$ and, since $u \in v \circ y$ we get $f(u) \geq f(v)$. Then we have $f(u) \geq f(a)$, and the proof is complete. □

Definition 2.5. An hypersemigroup H is called *regular* if for every $a \in H$ there exists $x \in H$ such that $a \in \{a\} * (x \circ a)$.

Proposition 2.6. *Let H be an hypersemigroup. The following are equivalent:*

(1) *H is regular.*

(1) *$a \in \{a\} * \{x\} * \{a\}$ for every $a \in H$.*

(2) *$A \subseteq H * A * H$ for every nonempty subset of H.*

Proposition 2.7. *Let H be a regular hypersemigroup and A an interior ideal of H. Then A is a subsemigroup of H.*

Proof. Since A is an interior ideal of H, we have $H * A * H \subseteq A$. Since H is regular, we have $A \subseteq A * H * A$. Then we have

$$\begin{aligned} A * A &\subseteq (A * H * A) * A = (A * H) * A * A \subseteq (H * H) * A * H \\ &\subseteq H * A * H \subseteq A, \end{aligned}$$

so A is a subsemigroup of H. □

Proposition 2.8. *Let H be a regular hypersemigroup and f a fuzzy interior ideal of H. Then f is a fuzzy ideal of H.*

Proof. Let $a, b \in H$ ($\Rightarrow f(a \circ b) \geq f(a)$ and $f(a \circ b) \geq f(b)$?)
Let $u \in a \circ b$. Then $f(u) \geq f(a)$. Indeed: Since $a \in H$ and H is regular, there exists $x \in H$ such that $a \in \{a\} * \{x\} * \{a\}$. Then

$$a \circ b \subseteq \{a\} * \{x\} * \{a\} * \{b\} = (a \circ x) * (a \circ b),$$

from which $u \in v \circ w$ for some $v \in a \circ x$, $w \in a \circ b$. We have $u \in v \circ w \subseteq \{v\} * (a \circ b)$ and $f\Big(\{v\} * (a \circ b)\Big) \geq f(a)$, thus we have $f(u) \geq f(a)$, and f is a fuzzy right ideal of H. We also have $f(u) \geq f(b)$. Indeed: Since $b \in H$ and H is regular, there exists $y \in H$ such that $b \in \{b\} * \{y\} * \{b\}$. Then we have

$$u \in a \circ b \subseteq \{a\} * \{b\} * \{y\} * \{b\} = (a \circ b) * (y \circ b).$$

Then $u \in s \circ t$ for some $s \in a \circ b$, $t \in y \circ b$. Then we have

$$u \in s \circ t \subseteq (a \circ b) * \{t\} = \{a\} * (b \circ t).$$

Since $f\Big(\{a\} * (b \circ t)\Big) \geq f(b)$, we obtain $f(u) \geq f(b)$, and f is a fuzzy left ideal of H. □

From the Propositions 2.4 and 2.8 we have the following

Theorem 2.9. *In regular hypersemigroups the concepts of fuzzy ideals and of fuzzy interior ideals coincide.*

Definition 2.10. An hypersemigroup H is called *intra-regular* if for every $a \in H$ there exist $x, y \in H$ such that $a \in (x \circ a) * (a \circ y)$.

Proposition 2.11. *Let H be an hypersemigroup. The following are equivalent:*

(1) *H is intra-regular.*

(2) *$a \in H * \{a\} * \{a\} * H$ for every $a \in H$.*

(3) *$A \subseteq H * A * A * H$ for every nonempty subset of H.*

Proposition 2.12. *Let H be an intra-regular hypersemigroup and A an interior ideal of H. Then A is a subsemigroup of H.*

Proof. Since A is an interior ideal of H, we have $H * A * H \subseteq A$. Since H is intra-regular, we have $A \subseteq H * A * A * H$. Then we have

$$\begin{aligned} A * A &\subseteq (H * A * A * H) * A = (H * A) * A * (H * A) \\ &\subseteq (H * H) * A * (H * H) \\ &\subseteq H * A * H \subseteq A. \end{aligned}$$

□

By Propositions 2.7 and 2.12, we have the following

Corollary 2.13. *In regular and in intra-regular hypersemigroups the interior ideals and the subidempotent interior ideals coincide.*

Proposition 2.14. *Let H be an intra-regular hypersemigroup and f is a fuzzy interior ideal of H. Then f is a fuzzy ideal of H.*

Proof. Let $a, b \in H$ and $u \in a \circ b$. Since $a \in H$ and H is intra-regular, there exist $x, y \in H$ such that $a \in \{x\} * \{a\} * \{a\} * \{y\}$. Then

$$a \circ b \subseteq \{x\} * \{a\} * \{a\} * \{y\} * \{b\} = (x \circ a) * \Big((a \circ y) * \{b\}\Big).$$

Then $u \in v \circ w$ for some $v \in x \circ a$, $w \in (a \circ y) * \{b\}$. We have

$$u \in v \circ w \subseteq (x \circ a) * \{w\}.$$

Since f is a fuzzy interior ideal of H, $f\Big((x \circ a) * \{w\}\Big) \geq f(a)$. Thus we get $f(u) \geq f(a)$, and f is a fuzzy right ideal of H. Since $b \in H$ and H is intra-regular, there exist $z, t \in H$ such that $b \in \{z\} * \{b\} * \{b\} * \{t\}$, then we have

$$a \circ b \subseteq \{a\} * \{z\} * \{b\} * \{b\} * \{t\} = \Big((a \circ z) * \{b\}\Big) * (b \circ t).$$

Then $u \in c \circ d$ for some $c \in (a \circ z) * \{b\}$, $d \in b \circ t$. Then $u \in c \circ d \subseteq \{c\} * (b \circ t)$ and $f\Big(\{c\} * (b \circ t)\Big) \geq f(b)$, so $f(u) \geq f(b)$, and f is a fuzzy right ideal of H. □
By Propositions 2.4 and 2.14, we have the following theorem

Theorem 2.15. *In intra-regular hypersemigroups the concepts of fuzzy ideals and fuzzy interior ideals coincide.*

Definition 2.16. An ideal A of an hypergroupoid H is called *proper* if $A \neq H$. An hypergroupoid H is called *simple* if does not contain proper ideals, that is, for every ideal A of H, we have $A = H$.

Lemma 2.17. *An hypersemigroup H is simple if and only if, for every $a \in H$, we have*

$$H = H * \{a\} * H.$$

Proof. $\Longrightarrow$. Let $a \in H$. The set $H * \{a\} * H$ is an ideal of H. Indeed, it is a nonempty subset of H, and we have

$H * (H * \{a\} * H) = (H * H) * \{a\} * H \subseteq H * \{a\} * H$ and

$(H * \{a\} * H) * H = H * \{a\} * (H * H) \subseteq H * \{a\} * H$.

Since H is simple, we have $H * \{a\} * H = H$.

$\Longleftarrow$. Let A be an ideal of H. Then $A = H$. Indeed: Take an element $x \in A$ $(A \neq \emptyset)$. By hypothesis, we have $H = H * \{x\} * H \subseteq (H * A) * H \subseteq A * H \subseteq A$, thus we have $H = A$. □

Definition 2.18. (cf. also [2]) An hypergroupoid H is called *fuzzy simple* if every fuzzy ideal of H is a constant function, that is, for every fuzzy ideal f of H and every $a, b \in H$, we have $f(a) = f(b)$.

Notation 2.19. Let H be an hypergroupoid and $a \in H$. We denote by I_a the subset of H defined as follows:

$$I_a = \{b \in H \mid f(b) \geq f(a)\}.$$

Proposition 2.20. *Let H be an hypergroupoid and f a fuzzy right ideal of H. Then the set I_a is a right ideal of H for every $a \in H$.*

Proof. Let $a \in H$. The set I_a is a right ideal of H. Indeed: Since $a \in I_a$, the set I_a is a nonempty subset of H. Moreover, $I_a * H \subseteq I_a$. Indeed: Let $x \in I_a * H$. Then $x \in H$ and $x \in u \circ v$ for some $u \in I_a$, $v \in H$. Since $x \in u \circ v$ and f is a fuzzy right ideal of H, we have $f(x) \geq f(u)$. Since $u \in I_a$, we have $f(u) \geq f(a)$. Then we have $f(x) \geq f(a)$. Since $x \in H$ and $f(x) \geq f(a)$, we have $x \in I_a$. Thus I_a is a right ideal of H. □

In a similar way we prove the following

Proposition 2.21. *Let H be an hypergroupoid and f a fuzzy left ideal of H. Then the set I_a is a left ideal of H for every $a \in H$.*

By Propositions 2.20 and 2.21 we have the following proposition

Proposition 2.22. *Let H be an hypergroupoid and f a fuzzy ideal of H. Then the set I_a is an ideal of H for every $a \in H$.*

Theorem 2.23. *An hypergroupoid H is simple if and only if it is fuzzy simple.*

Proof. $\Longrightarrow$. Let f be a fuzzy ideal of H and $a, b \in H$. Then $f(a) = f(b)$. Indeed: Since f is a fuzzy ideal of H and $a \in H$, by Proposition 2.22, the set $I_a = \{b \in H \mid f(b) \geq f(a)\}$ is an ideal of H. Since H is simple, we have $I_a = H$. Then $b \in I_a$, so $f(b) \geq f(a)$. By symmetry, we get $f(a) \geq f(b)$, thus $f(a) = f(b)$. $\Longleftarrow$. Let H be fuzzy simple and I an ideal of H. Then $I = H$. Indeed: Let $x \in H$. Since I is an ideal of H, the characteristic function f_I is a fuzzy ideal of H. Since H is fuzzy simple, f_I is a constant function, that is, $f_I(y) = f_I(z)$ for every $y, z \in H$. Take an element $a \in I$ ($I \neq \emptyset$). Then we have $f_I(x) = f_I(a) = 1$, so $x \in I$. Thus H is simple. □

Theorem 2.24. *Let H be an hypersemigroup. Then H is simple if and only if every fuzzy interior ideal of H is a constant function.*

Proof. $\Longrightarrow$. Let f be a fuzzy interior ideal of H and $a, b \in H$. Then $f(a) = f(b)$. Indeed: Since $a \in H$ and H is simple, by Lemma 2.17, we have $H = H * \{a\} * H$.

Since $b \in H$, we have $b \in (x \circ a) * \{y\}$ for some $x, y \in H$. Since f is a fuzzy interior ideal of H, we have $f(b) \geq f(a)$. By symmetry, we get $f(a) \geq f(b)$, so $f(a) = f(b)$.

$\Longleftarrow$. Let f is a fuzzy ideal of H. By Proposition 2.4, f is a fuzzy interior ideal of H. By hypothesis, f is a constant function. Thus H is fuzzy simple. Then, by Theorem 2.23, H is simple. □

As a consequence we have the following:

Theorem 2.25. *If H is an hypersemigroup H, then the following are equivalent:*

(1) *H is simple.*

(2) *$H = H * \{a\} * H$.*

(3) *H is fuzzy simple.*

(4) *Every fuzzy interior ideal of H is a constant function.*

References

[1] N. Kehayopulu, M. Tsingelis, Fuzzy interior ideals in ordered semigroups, Lobachevskii J. Math. **21** (2006), 65–71.

[2] N. Kuroki, Fuzzy semiprime ideals in semigroups, Fuzzy Sets and Systems **8** (1982), 71–79.

University of Athens
Department of Mathematics
15784 Panepistimiopolis, Greece
email: nkehayop@math.uoa.gr